arXiv:1604.05755v1 [math.GR] 19 Apr 2016

Algebras of conjugacy classes in symmetric groups

Yu.A.Neretin[1]

In 1999 V. Ivanov and S. Kerov observed that structure constants of algebras of conjugacy classes of symmetric groups S_n admit a stabiluzation (in a nontrivial sence) as $n \to \infty$. We extend their construction to a certain class of pairs of groups $G \supset K$ and algebras of conjugacy classes of G with respect to K (as $S_{kn} \supset S_n$, $S_n \times S_n \supset S_n$, etc.).

1 The statement of the paper

1.1. Notation. Denote by $\mathbb{Z}_+$ the set $\{0, 1, 2, \dots\}$. Denote by J_n the subset $\{1, 2, \dots, n\}$ in $\mathbb{N}$. Let Y be a finite or countable set. Denote by $S(Y)$ the group of all finitely supported permutations of Y. By S_n we denote the group of permutations of J_n, by S_∞ the group $S(\mathbb{N})$.

Let G be a finite group, K a subgroup. By $G//K$ we denote the set of conjugacy classes of G with respect to K, by $\mathbb{C}(G//K)$ the convolution algebra of functions on G invariant with respect to conjugations by elements of K,

$$f(\sigma g \sigma^{-1}) = f(g) \qquad \text{for } \sigma \in K.$$

By $\mathbb{C}(G)$ we denote the group algebra of G. The symbol $*$ denotes the multiplication in $\mathbb{C}(G)$ (the convolution).

1.2. Groups G_n. Fix a finite set I. Consider a finite or countable set[2] X, the product $I \times \mathbb{N}$ and the unions

$$V = X \cup (I \times \mathbb{N}), \qquad V_n = X \cup (I \times J_n) \subset V,$$

see Fig. 1. The group S_∞ acts on $\mathbb{N}$, therefore it acts on the product $I \times \mathbb{N}$. We can imagine $I \times \mathbb{N}$ as a table with $\#I$ infinite rows. The group S_∞ acts by permutations of columns.

Next, we consider the trivial action of S_∞ on X, this determines an action of S_∞ the whole V.

Consider a subgroup $G \subset S(V)$ containing S_∞. Set

$$G_n := G \cap S(V_n).$$

In other words, we consider elements of G supported by V_n.

Examples. a) $X = \varnothing$, $G = S(I \times \mathbb{N})$.

b) $X = \varnothing$, $G = S_\infty \times \cdots \times S_\infty$, its subgroup S_∞ is the diagonal.

c) X is a finite set, $\#I = 1$, $G = S(X \cup \mathbb{N})$.

[1]Supported by the grant FWF, P28421.

[2]The case $X = \varnothing$ is sufficiently interesting.



Figure 1: Sets $V \supset V_n$.

d) $X = \varnothing$, H is a group of permutations of I, G is the wreath product of S_∞ and H.

1.3. Algebras $\mathbb{C}(G_n//S_n)$. Let $\overline{\overline{g}}$, $\overline{\overline{h}}$, $\overline{\overline{r}}$, ... range in $\coprod_{j=0}^{\infty} G_j//S_j$. Let denote by g, h, r their representatives.

Fix $n \in \mathbb{Z}_+$, $\overline{\overline{g}} \in G_n//S_n$, $g \in \overline{\overline{g}}$, and $N \geqslant n$. For any n-element set $\Omega \subset \mathbb{N}$ we define the sum

$$R(g, \Omega) := \sum_{\sigma : J_n \to \Omega} \sigma g \sigma^{-1} \quad \in \mathbb{C}(G),$$

where σ ranges in the set of bijective maps $J_n \to \Omega$. Explain this more carefully. Any σ determines a bijection $I \times J_n \to I \times \Omega$, we denote it by the same symbol σ. Thus, $\sigma g \sigma^{-1}$ is a map

$$\sigma g \sigma^{-1} : X \cup (I \times \Omega) \to X \cup (I \times \Omega).$$

We extend it to a map $V_N \to V_N$ in a trivial way,

$$\sigma g \sigma^{-1}(w) := w, \qquad \text{if } w \in I \times (J_n \setminus \Omega.)$$

Next, we consider the sum

$$A_N[\overline{\overline{g}}] = \sum_{\Omega \subset J_N :\ \#\Omega = n} R(g, \Omega).$$

For $N < n$, we set $A_N[\overline{\overline{g}}] = 0$.

Let $n < m \leqslant N$. For $g \in G_n$ denote by g_n^m the same permutation regarded as an element of G_m. Then

$$A_N[\overline{\overline{g_n^m}}] = \frac{(N-n)!}{(N-m)!} A_N[\overline{\overline{g}}].$$

There are no other linear dependences between elements $A_N[\overline{\overline{g}}]$.

Theorem 1.1 a) *Let $\overline{\overline{g}}$, $\overline{\overline{h}}$, $\overline{\overline{r}}$ range in the disjoint union $\coprod_{j=0}^{\infty} G_j//S_j$. Then there are non-negative integers $a^{\overline{\overline{r}}}_{\overline{\overline{g}}, \overline{\overline{h}}}$, which do not depend on N, such that for each N*

$$A_N[\overline{\overline{g}}] * A_N[\overline{\overline{h}}] = \sum_{\overline{\overline{r}}} a^{\overline{\overline{r}}}_{\overline{\overline{g}}, \overline{\overline{h}}} A_N[\overline{\overline{r}}]. \tag{1.1}$$

b) *Consider a linear space* $\mathcal{B}[G]$ *with a basis consisting of symbols* $A[\overline{\overline{g}}]$, *where* $\overline{\overline{g}}$ *ranges in* $\coprod_{j=0}^{\infty} G_j//S_j$. *Then the formula*

$$A[\overline{\overline{g}}] * A[\overline{\overline{h}}] = \sum_{\overline{\overline{r}}} a^{\overline{\overline{r}}}_{\overline{\overline{g}},\overline{\overline{h}}} A[\overline{\overline{r}}], \tag{1.2}$$

determines a structure of an assosiative algebra on $\mathcal{B}[G]$.

A proof of the theorem is contained in Section 2. The algebra $\mathcal{B}[G]$ is a more handable object that it seems at the first glance, this is discussed in Section 3.

2 Proof

2.1. Partial bijections. We say that a *partial bijection* λ of a set Y to a set Z is a bijection of a subset $A \subset Y$ to a subset $B \subset Y$ (we admit the case $A = B = \varnothing$). We define

$$\operatorname{rk}\lambda := \#A = \#B, \qquad \operatorname{dom}\lambda := A, \qquad \operatorname{im}\lambda := B.$$

Denote by $\mathrm{PB}(Y, Z)$ the set of all partial bijections $Y \to Z$.

For partial bijections $\mu : W \to Y$, $\lambda : Y \to Z$ we define their product $\lambda\mu : W \to Z$. We say that $w \in \operatorname{dom}\lambda\mu$ if $w \in \operatorname{dom}\mu$ and $\mu(w) \in \operatorname{dom}\lambda$. In this case we set $\lambda\mu(w) = \lambda(\mu(w))$

2.2. Local bijections. Following Ivanov and Kerov [1] we define a semigroup of local bijections. Let Y be a finite or countable set. A *local bijection* ω is a bijection from a finite subset $\Omega \subset Y$ to Ω. We denote such local bijection by $(\!(\omega, \Omega)\!)$. Any local bijection admits a canonical extension to an element $\widetilde{\omega} \in S(Y)$, we set $\widetilde{\omega}y = y$ if $y \notin \Omega$. We define the product of local bijections by

$$(\!(\omega, \Omega)\!) \circ (\!(\mu, M)\!) = \left(\!\!\left(\widetilde{\omega} \cdot \widetilde{\mu}\Big|_{\Omega \cup M}, \Omega \cup M\right)\!\!\right).$$

Denote by $\mathcal{L}(Y)$ the semigroup of all local bijections of Y.

Remark. A local bijection determines a partial bijection. But the $\circ$-product differs from the product of partial bijections. ⊠

The group $S(Y)$ acts on $\mathcal{L}(Y)$ by conjugations in the obvious way.

On the other hand, we have a natural forgetting homomorphism $\iota : \mathcal{L}(Y) \to S(Y)$ defined by

$$\iota(\!(\omega, \Omega)\!) = \widetilde{\omega}.$$

2.3. The semigroup algebra for local bijections. Denote by $\mathbb{C}[\mathcal{L}(Y)]$ the space of all formal series of the form

$$\sum_{(\!(\omega,\Omega)\!) \in \mathcal{L}(Y)} a_{(\!(\omega,\Omega)\!)} (\!(\omega, \Omega)\!), \qquad a_{(\!(\omega,\Omega)\!)} \in \mathbb{C}.$$

This space is equipped with a convolution

$$\sum_{((\omega,\Omega))} a_{((\omega,\Omega))}\,((\omega,\Omega)) * \sum_{((\mu,M))} b_{((\mu,M))}\,((\mu,M)) = \sum_{((\nu,N))} c_{((\nu,N))}\,((\nu,N)),$$

where

$$c_{((\nu,N))} = \sum_{((\omega\Omega)),((\mu M)):\ \Omega\cup M=N,\ \widetilde{\omega}\widetilde{\mu}=\widetilde{\nu}} a_{((\omega,\Omega))} b_{((\mu,M))}.$$

(this sum is finite).

For any subset Z in Y we have a homomorphism $\pi_Z^Y : \mathbb{C}[\mathcal{L}(Y)] \to \mathbb{C}[\mathcal{L}(Z)]$ defined on generators by

$$\pi((\omega,\Omega)) = \begin{cases} ((\omega,\Omega)), & \text{if } \Omega \subset Z \\ 0, & \text{otherwise.} \end{cases}$$

For a finite Y we have a homomorphism of algebras $\iota : \mathbb{C}[\mathcal{L}(Y)] \to \mathbb{C}[S(Y)]$ defined by

$$\iota \sum_{((\omega,\Omega))} a_{((\omega,\Omega))}\,((\omega,\Omega)) = \sum_{((\omega,\Omega))} a_{((\omega,\Omega))}\widetilde{\omega} = \sum_{g\in S(Y)} \Big[\sum_{\Omega:\, g\Omega=\Omega} a_{((g\big|_\Omega,\Omega))} \Big] g$$

(for an infinite set Y the sum in square brackets is infinite).

2.4. Elements $B[g]$. Let $\overline{\overline{g}} \in G_n//S_n$, $g \in \overline{\overline{g}}$. For any an n-element subset $\Omega \subset \mathbb{N}$ we define an element of $\mathbb{C}[\mathcal{L}(V)]$ by

$$\widetilde{R}(\overline{\overline{g}},\Omega) = \sum_{\sigma: J_n\to\Omega} \big((\sigma g\sigma^{-1}, X\cup(I\times\Omega)\big)\big),$$

where the summation is taken over all bijections $\sigma : J_n \to \Omega$.

Next, we write

$$B[\overline{\overline{g}}] := \sum_{\Omega:\,\#\Omega=n} \widetilde{R}(\overline{\overline{g}},\Omega).$$

By construction these elements are invariant with respect to conjugations by elements of S_∞,

For any $\overline{\overline{g}} \in G_n//S_n$, $\overline{\overline{h}} \in G_m//S_m$ the convolution $B[\overline{\overline{g}}] * B[\overline{\overline{h}}]$ is S_∞-invariant, therefore it has the form

$$B[\overline{\overline{g}}] * B[\overline{\overline{h}}] := \sum_{\overline{\overline{r}}\in\coprod G_k//S_k} a^{\overline{\overline{r}}}_{\overline{\overline{g}},\overline{\overline{h}}} B[\overline{\overline{r}}].$$

2.5. Proof of the theorem. First, we apply the map π corresponding the set V and its subset V_N. For any $\overline{\overline{g}}$ and $n \in \mathbb{N}$ we get an element

$$B_N[\overline{\overline{g}}] := \pi B[\overline{\overline{g}}].$$

Since π is a homomorphism of algebras, we get

$$B_N[\overline{\overline{g}}] * B_N[\overline{\overline{h}}] := \sum_{r \in \coprod_{k=0}^N G_k//S_k} a^{\overline{\overline{r}}}_{\overline{\overline{g}}, \overline{\overline{h}}} B_N[\overline{\overline{r}}],$$

where g, h range in $\coprod_{k=0}^N G_k//S_k$.

Next, we apply the forgetting map $\mathbb{C}[\mathcal{L}(V_n)] \to \mathbb{C}[S(V_n)]$ to elements $B_N[\overline{\overline{g}}]$. Evidentely, we have

$$\iota(B_N[\overline{\overline{g}}]) = A_N[\overline{\overline{g}}].$$

This implies our statement.

3 Remarks

3.1. An expression for products. Let $g \in G_n$, $h \in G_k$. Let λ ranges in a partial bijections $J_k \to J_n$. Fix λ, denote $d = \operatorname{rk}\lambda$. Fix any pair of injective maps $\sigma : J_n \to J_{n+k-d}$, $\tau : J_k \to J_{n+k-d}$ such that the product $\sigma^{-1}\tau$ in the category of partial bijections is λ. Define the conjugacy class

$$\overline{\overline{g \circledast_\lambda h}} := \overline{\overline{\sigma(\!(g, V_n)\!)\sigma^{-1} \circ \tau(\!(h, V_k)\!)\tau^{-1}}} \in G_{n+k-d}//S_{n+k-d},$$

it does not depend on the choice of σ and τ.

Proposition 3.1

$$B[\overline{\overline{g}}] * B[\overline{\overline{h}}] = \sum_{\lambda \in \mathrm{PB}(J_k, J_n)} B[\overline{\overline{g \circledast_\lambda h}}]. \tag{3.1}$$

Proof. We expand the product $B[\overline{\overline{g}}] * B[\overline{\overline{h}}]$ by definition as a sum

$$\sum_{\sigma,\tau} \sigma(\!(g, V_n)\!)\sigma^{-1} \circ \tau(\!(h, V_k)\!)\tau^{-1} = \sum_{\tau,\sigma} (\!(\sigma g \sigma^{-1}, \sigma^{-1} V_n)\!) \circ (\!(\tau h \tau^{-1}, \tau^{-1} V_k)\!), \tag{3.2}$$

where $\sigma : J_n \to \mathbb{N}$, $\tau : J_k \to \mathbb{N}$ are injective maps. We wish to show that this expansion coincides with (3.1). It is sufficient to identify the following subsum of (3.2)

$$\sum_{\sigma,\tau:\ \sigma J_n \cup \tau J_k = J_{n+k-d}} (\!(\sigma g \sigma^{-1}, \sigma^{-1} V_n)\!) \circ (\!(\tau h \tau^{-1}, \tau^{-1} V_k)\!) \tag{3.3}$$

and the subsum of (3.1) consisting of all summands of the form

$$(\!(r, V_{n+k-d})\!).$$

Notice that the group S_{n+k-d} acts on the set of summands of (3.3) by

$$u :\ (\sigma, \tau) \mapsto (u\sigma, u\tau).$$



Figure 2: The left part. A product $\sigma(\!(g, V_n)\!)\,\sigma^{-1} \circ \tau(\!(h, V_k)\!)\,\tau^{-1}$. Here $k = 5$, $n = 7$, $d = 3$, $n + k - d = 9$. We draw 7 copies of the set J_{n-k+1}. Maps τ^{-1}, τ, σ^{-1}, σ are shown by arcs. We mark $\operatorname{dom}\lambda$, $\operatorname{im}\lambda$, and $\sigma \operatorname{dom}\lambda = \tau^{-1} \operatorname{im}\lambda$ as fat points. Arcs connecting fat points are thick. Boxes correspond to elements $(\!(h, V_k)\!) \in G_k$ and $(\!(g, V_n)\!) \in G_n$.
The right part. A canonical form of the pair (τ, σ).



Figure 3: A piece of a well-equipped surface.

We refer to Fig. 2. This action corresponds to simultanious application of a substitution u to rows number 1, 4, 7. Rows 2, 3, 5, 6 remain to be fixed, and arcs move corresponding permutations of their ends. We can put fat points to positions 1, 2, ..., d. Moreover, we can make $\tau^{-1}u$ monotone on J_d:

$$i < j \leqslant d \qquad \Rightarrow \qquad \tau^{-1}u(i) < \tau^{-1}u(j).$$

(on Fig. 2 this means that the correspondimg arcs have no intersections).

Next, we can put points of $\tau J_k \setminus \tau J_d$ to points of $J_k \setminus J_d$. Moreover, we can make $\tau^{-1}u$ monotone on $J_k \setminus J_d$. Finally, we can put points of $\sigma J_n \setminus \sigma J_d$ to points of $J_{n+k-d} \setminus J_k$. Moreover, we can make $\sigma^{-1}u$ monotone on $J_{n+k-d} \setminus J_k$.

This determines u in a unique way. On the other hand, the partial bijection $\sigma^{-1}\tau$ does not change under this tranformation, and the new pair $(u\sigma, u\tau)$ is uniquely determined by $\operatorname{dom}\sigma^{-1}\tau$, $\operatorname{im}\sigma^{-1}\tau$ and the map $\sigma^{-1}u : J_d \to \operatorname{im}\sigma^{-1}\tau$.

Thus, we see that orbits of the group S_{n+k-d} on the set of summands of (3.3) are enumerated by partial bijections λ. $\square$

3.2. Example. Consider the group $G = S_\infty \times S_\infty$ and conjugacy classes with respect to its diagonal subgroup S_∞. Then $G_N = S_N \times S_N$. Conjugacy classes with respect to the diagonal S_N can be described in the following language (see [3]).

We say that a triangulated two-dimensional oriented closed surfaceis *well-equipped* (see Fig. 3), if

— for earh face we assign a sign (+) or (−) in a checker order;

— edges are colored by red, yellow, blue; colors of edges of each face are different; in plus-triangles these colors are located clockwise, on minus-triangles anti-clockwise.

We admit non-connected surfaces. Also, we admit pairs of triangles, which have two or three common edges as on Fig. 4. Surfaces are defined up to the combinatorial equivalence.

Denote by Ξ_N the set of all equpped surfaces with $2N$ faces. For any element $\mathcal{R} \in \Xi_N$ we can asign a conjugacy class $\in G_N//S_N$. Denote by $\mathcal{R}_+$ the set of plus-faces of $\mathcal{R}$, by $\mathcal{R}_-$ the set of minus-faces. Any red edge has a plus-side and



Figure 4: Surfaces $\mathcal{R}$, $\mathcal{Q}$ and a partial bijection $\mathcal{R}_-$ to $\mathcal{Q}_+$.

minus side. This determines a bijection $\sigma_{red} : \mathcal{R}_+ \to \mathcal{R}_-$. In the same way, we define bijections σ_{yellow}, $\sigma_{blue} : \mathcal{R}_+ \to \mathcal{R}_-$. Therefore, we have bijections

$$\sigma_{yellow}^{-1}\sigma_{red},\ \sigma_{yellow}^{-1}\sigma_{blue} : \mathcal{R}_+ \to \mathcal{R}_+.$$

Since we can enumerate elements of $\mathcal{R}_+$ in an arbitrary way, we get a congugacy class $\in G_N//S_N$.

Such objects were a topic of [3]. It is easy to describe the algebra $\mathcal{B}[G]$ in this case. A basis of $\mathcal{B}[G]$ is enumerated by elements of the set $\coprod_{n=0}^{\infty}\Xi_n$. Let $\mathcal{R} \in \Xi_n$, $\mathcal{Q} \in \Xi_k$. Let $\lambda : \mathcal{R}_- \to \mathcal{Q}_+$ be a partial bijection (see Fig. 4). Consider the disjoint union $\mathcal{R}\coprod\mathcal{Q}$ and let us perform the following transformations. For each face $A \in \operatorname{dom}\lambda$, we take the face $\lambda(A) \in \operatorname{im}\lambda$, remove both faces and identify their boundaries according colors of edges. After this, we get a compact two-dimensional simplicial complex, and each edge of the complex is contained in precisely two faces. In fact, this is a surface, but some vertices of the surface can be glued one with another (see Fig. 5). Cutting all such gluengs we get a new surface

$$\mathcal{R} \circledast_\lambda \mathcal{Q}.$$

In this notation,

$$B[\mathcal{R}] * B[\mathcal{Q}] = \sum_{\lambda\in \mathrm{PB}(\mathcal{R}_-,\mathcal{Q}_+)} B[\mathcal{R} \circledast_\lambda \mathcal{Q}].$$

3.3. The involution. The map $g \mapsto g^{-1}$ determines an involution in the algebra $\mathcal{L}[V]$ and involution in $\mathcal{B}[G]$,

$$B[\overline{\overline{g}}]^\star := B[\overline{\overline{g^{-1}}}].$$

Evidentely,

$$(B[\overline{\overline{g}}] * B[\overline{\overline{h}}])^\star = B[\overline{\overline{h}}]^\star * B[\overline{\overline{g}}]^\star.$$

3.4. The filtration. Let $\mathcal{B}[G]_n$ be the subspace in $\mathcal{B}[G]$ generated by all

$$B[\overline{\overline{g}}], \qquad \text{where } g \text{ ranges in } \coprod_{j=0}^{n} G_j//S_j.$$

We get an increasing filtration,

$$\dots \subset \mathcal{B}[G]_n \subset \mathcal{B}[G]_{n+1} \subset \dots$$

Evidentely,

$$V \in \mathcal{B}[G]_k, \quad W \in \mathcal{B}[G]_l \quad \Rightarrow \quad V * W \in \mathcal{B}[G]_{k+l}.$$

3.5. The associated graded algebra. We construct the graded algebra $gr\mathcal{B}[G]$ in the usual way. Namely, the product

$$\mathcal{B}[G]_k \times \mathcal{B}[G]_k \ \to \mathcal{B}[G]_{k+l}$$



Figure 5: A surface whose vertices are glued.

determines a map

$$\mathcal{B}[G]_k/\mathcal{B}[G]_{k-1} \times \mathcal{B}[G]_l/\mathcal{B}[G]_{l-1} \ \to \ \mathcal{B}[G]_{k+l}/\mathcal{B}[G]_{k+l-1}.$$

In this way, we get a structure of an associative algebra on

$$\mathrm{gr}\mathcal{B}[G] = \bigoplus_{k=0}^{\infty} \mathcal{B}[G]_k/\mathcal{B}[G]_{k-1}.$$

Each subspace $\mathcal{B}[G]_k/\mathcal{B}[G]_{k-1}$ has a natural basis enumerated by elements of $G_k//S_k$.

It is easy to describe the multiplication in $\mathrm{gr}\mathcal{B}[G]$: in the sum in the right-hand side of (3.1) we leave only the first summand (corresponding to the parial bijection of rank 0).

Formulate this more precisely. For $n,\ k \in \mathbb{Z}_+$ denote by $\theta_{n,k}$ the partial bijection $J_n \to J_{n+k} \setminus J_k$ defined by $j \mapsto j+k$.

Proposition 3.2 a) *For* $\overline{\overline{g}} \in G_n//S_n$, $\overline{\overline{h}} \in G_k//S_k$, *their product in* $\mathrm{gr}\mathcal{B}[G]$ *is*

$$B[\overline{\overline{g}}] \diamond B[\overline{\overline{h}}] = B[\overline{\overline{\theta_{n,k}(\!(g, V_n)\!)\theta_{n,k}^{-1} \circ (\!(h, V_k)\!)}}].$$

b) *If* $X = \varnothing$, *then the algebra* $\mathrm{gr}\mathcal{B}[G]$ *is commutative.*



Figure 6: The $\bullet$-multiplication.

In fact, we get a semigroup structure on $\coprod_{j=0}^{\infty} G_j//S_j$ given by

$$\overline{\overline{g}} \bullet \overline{\overline{h}} := \overline{\overline{\theta_{n,k}(\!(g, V_n)\!)\theta_{n,k}^{-1} \circ (\!(h, V_k)\!)}}$$

(see Fig.6). The algebra $gr\mathcal{B}[G]$ is the semigroup algebra of this semigroup (so it is similar to algebraic structures, which were considered in [3] and [2], Chapter VIII).

Example. For $G = S_\infty \times S_\infty$ the $\bullet$-product corresponds to the disjoint union of well-equipped surfaces.

3.6. The Poisson bracket. Let $X = \varnothing$. Consider the map

$$\mathcal{B}[G]_k \times \mathcal{B}[G]_l \ \to \ \mathcal{B}[G]_{k+l-1}$$

given by

$$(V, W) \mapsto [V, W] = V * W - W * V.$$

As usual, we get a map

$$\mathcal{B}[G]_k/\mathcal{B}[G]_{k-1} \times \mathcal{B}[G]_l/\mathcal{B}[G]_{l-1} \ \to \ \mathcal{B}[G]_{k+l-1}/\mathcal{B}[G]_{k+l-2}$$

and a structure of a Lie algebra on the space $gr\mathcal{B}[G]$. It is easy to write a formula for the bracket

$$\big[B[\overline{\overline{g}}], B[\overline{\overline{h}}]\big]_{gr} = \sum_{\lambda \in \mathrm{PB}(J_k, J_n)\colon \mathrm{rk}\,\lambda = 1} \Big(B[\overline{\overline{g \circledast_\lambda h}}] - B[\overline{\overline{h \circledast_{\lambda^{-1}} g}}] \Big).$$

Of cource, a partial bijection $J_k \to J_n$ of rank 1 is a pair $\alpha \in J_k$, $\beta \in J_n$.

References

[1] Ivanov V., Kerov S., *The Algebra of Conjugacy Classes in Symmetric Groups and Partial Permutations*, Zapiski nauchn.semin. POMI RAN, 256 (1999), 95–120; J. of Math. Sci. (New York), 2001, 107:5, 4212-4230.

[2] Neretin Yu.A. *Categories of symmetries and infinite-dimensional groups.* Oxford University Press, 1996.

[3] Neretin Yu.A., *Infinite symmetric groups and combinatorial constructions of topological field theory type.* Russian Mathematical Surveys, 2015, 70:4, 715-773.

Math.Dept., University of Vienna,
Oskar-Morgenstern-Platz 1, 1090 Wien;
& Institute for Theoretical and Experimental Physics (Moscow);
& Mech.Math.Dept., Moscow State University;
Institute for information transmission problems (Moscow);
e-mail: neretin(at) mccme.ru
URL:www.mat.univie.ac.at/∼neretin